

OPIC
OFFICE DE LA PROPRIÉTÉ
INTELLECTUELLE DU CANADA



CIPO
CANADIAN INTELLECTUAL
PROPERTY OFFICE

CHINA HUI KANG

(21) (A1)	2,180,193
(22)	1996/06/28
(43)	1996/12/29

(51) Int.Cl. ⁶ A61K 38/21; A61K 38/20; A61K 31/135

(19) (CA) APPLICATION FOR CANADIAN PATENT (12)

(54) Enhancing Host Immunity Against Viral Infections

(72) Baral, Edward - Canada ;
Nagy, Eva - Canada ;
Dawood, Magdy - Canada ;
Berczi, Istvan - Canada ;

(71) UNIVERSITY OF MANITOBA - Canada ;
MANITOBA CANCER TREATMENT AND RESEARCH FOUNDATION -
Canada ;

(30) (US) 60/000,576 1995/06/28

(57) 26 Claims

Notice: This application is as filed and may therefore contain an
incomplete specification.



Industrie Canada Industry Canada

OPIC CIPO

Canada

BAD ORIGINAL

ABSTRACT OF THE DISCLOSURE

Cell mediated immunity against persistent viral infection in a host is potentiated by administering to the host an effective amount of at least one antiestrogen capable of enhancing cytotoxic T lymphocytes in the host, such as a triphenylethylene tamoxifen, particularly tamoxifen or toremifene. This administration leads to the elimination of all infected cells and to the complete clearance of virus and virus induced tumor cells from the body, leaving protective cell mediated immunity. This therapy requires MHC restricted virus specific CTLs which then will kill virus infected very specifically, which may be initially present or formed in vivo or in vitro. The effect of the antiestrogen may be enhanced by the use of interferons optionally with indomethacin.

BAD ORIGINAL

1887-168 MIS 739 1996 06 21 D1

TITLE OF INVENTIONENHANCING HOST IMMUNITY AGAINST VIRAL INFECTIONSFIELD OF INVENTION

The present invention relates to immunology.

BACKGROUND OF INVENTION

5 Viruses carry infectious genetic material (either DNA or RNA) with the capacity of entering susceptible host cells and redirecting the biosynthetic machinery of the infected cells towards the multiplication of viral particles (virions). The profound conversion of the host
10 cell toward viral production results in cell lysis in most cases (cytotoxic viruses), whereas in other cases the virus has no cytolytic effect, but rather persists within infected cells in the absence of morphological alteration. However, such virus infected cells may show
15 a severe impairment of functions that are not necessary for cell survival (e.g. the secretion of hormones, antibodies, and other important molecules) (Fenner & White, 1976; de la Torre et al., 1991).

The DNA viruses most frequently causing persistent
20 infections in man are cytomegaloviruses, the Epstein-Barr virus, herpes viruses (types I and II), hepatitis viruses (B, D), papilloma viruses, and occasionally measles virus. Recently, an association has been found between type II diabetes and the persistence of cytomegalovirus
25 in the pancreas. Herpes viruses persist in nerve cells and may cause severe inflammatory disease in the nervous system and also in various other parts of the body after having travelled along the nerve fibers and triggering inflammation in the area of innervation of the fibers
30 infected. The Epstein-Barr virus is known to cause infectious mononucleosis, and contributes to the pathogenesis of Burkitt's lymphoma, primarily in Africa, and nasopharyngeal carcinoma in the Orient. Hepatitis viruses cause inflammatory liver disease which may become

BAD ORIGINAL

5 persistent in some individuals leading to cirrhosis. In
addition, the hepatitis B virus has been identified as a
contributing agent to the genesis of hepatocellular
carcinomas. The measles virus may also persist in the
nervous system and is known to cause the condition called
subacute sclerosing panencephalitis. Papilloma viruses
(Papova group) exist in many species, including man, and
are known to cause common warts and genital tumors.
Progressive multifocal encephalopathy is also caused by
10 Papova virus (Fenner & White, 1976; Kimelman et al.,
1985; Chow, 1993).

The common RNA viruses with the potential of causing
persistent infections in man are hepatitis A and C,
rubella, the human T lymphocyte virus, types I and II
15 (HTLV), which are known to cause lymphomas, and the human
immunodeficiency virus (HIV) which is recognized as the
causative agent of acquired immunodeficiency syndrome
(AIDS). HTLV and HIV possess an enzyme, called reverse
transcriptase, which is capable of transcribing the viral
20 RNA to a DNA template which in turn will permanently be
inserted in the genome of the host cell. For this reason
they are classified as retroviruses. Both DNA and RNA
tumor viruses carry genes that are capable of growth
stimulation of the host cell (viral oncogenes) (Fenner &
25 White, 1976; Chow, 1993).

Persistent viral infections of farm animals that
have economic significance are: avian leukosis caused by
retroviruses (Okazaki et al., 1982); bovine leukosis,
another retroviral disease (Daniel et al., 1993); bovine
30 viral diarrhea (RNA pestivirus) (Thiel et al., 1993);
bluetongue, a disease of ruminants transmitted by insects
and caused by RNA viruses (arboviruses) (Sellers, 1981);
equine infectious anemia caused by retroviruses (Sellon
et al., 1994; Kennedy, 1992); retroviral disease of sheep
35 caused by the Maedi-Visna virus (Kennedy, 1992); porcine

BAD ORIGINAL

reproductive and respiratory syndrome caused by RNA lentivirus (Thiel et al., 1993).

Virus infected cells and virus induced tumors carry specific antigens on their surface which are recognized
5 by the immune system of the host. These antigens may stimulate cell mediated and humoral immune reactions. Major histocompatibility complex antigen (MHC) restricted cytotoxic T lymphocytes, which are capable of
10 specifically killing virus infected cells, are fundamental to host defence against viruses. However, it is recognized that natural killer cells and activated macrophages and the cytokines that are capable of
15 activating these effector cells, namely IL-2, interferon- α and γ , and migration inhibitory factor, also contribute significantly to host defence against viral infections. Interferons α and γ have the capacity to inhibit the
intracellular replication of viruses. Although some antibodies are capable of neutralizing viral particles,
humoral immunity is considered to play a secondary role
20 in host defence against viral infections (Fenner & White, 1976; Rinaldo & Torpey, 1993).

Initially, cell mediated immunity is present in immunocompetent hosts infected with persistent or
oncogenic viruses. However, with time a gradual
25 reduction occurs in the cell mediated antiviral/antitumor response, which is followed by nonprotective humoral responses, and some of the antibodies produced may actually interfere with cell mediated immunity (enhancing
antibodies). Virus infected non-neoplastic cells may be
30 similarly protected. In the case of tumors, immune derived growth factors may actually have a direct promoting effect on tumor growth which is known as immunostimulation. Apart from inducing a switch in the
host immune response, which permits the persistence of
35 viral infections and the growth of virus induced tumors, numerous other mechanisms have been proposed by which

immunodestruction of infected cells can be avoided. Some of these are: molecular mimicry, growing in immunologically privileged sites (such as the central nervous system), rendering immunocytes nonfunctional by direct infection, triggering nonspecific cytotoxic and autoimmune reactions (Oldstone & Rall, 1993; Tishon et al., 1993; Roger et al., 1994; Seligmann, 1990). HIV exposed individuals who are capable of resisting the establishment of persistent infection possess MHC-restricted cytotoxic T lymphocytes, but remain seronegative (Rowland-Jones et al., 1995).

Recent evidence indicates that virus specific CTL can be recovered from HIV infected individuals and from AIDS patients with active disease by in vitro stimulation with antigen and cytokines such as interleukin-2 or interleukin-12 (Clerici et al., 1993; Walker, 1989; Whiteside, 1993). Similarly, progressing neoplasms contain tumor infiltrating lymphocytes (TIL) which are inefficient in rejecting the tumor in vivo, but can be reactivated in vitro to exert tumor specific cytotoxicity (Berczi et al., 1973; Schendel & Gansbacher, 1993; Ioannides et al., 1993). Such reactivated CTL is being tested at present for the immunotherapy of cancer and of some viral infections (Kradin et al., 1989; Topalian et al., 1988; Torpey et al., 1993; Whiteside et al., 1993; Riddell et al., 1993).

SUMMARY OF THE INVENTION

We have discovered that the treatment of Epstein-Barr virus (EBV) transformed human B lymphocytes with the antiestrogenic agents, tamoxifen (TX) or toremifene (TO), enhances the destruction of these cells when used as targets for specific autologous CTL and LAK cells (see Tables 1 and 2 below). Currently, tamoxifen and toremifene are used in medicine for the treatment of estrogen receptor positive breast carcinomas and some

BAD ORIGINAL

other estrogen dependent tumors. Human B lymphocytes do not express classical estrogen receptors, which suggests that the sensitizing effect is mediated by some other receptor.

5 Having regard to the new observations we have made, the present invention, in one aspect, provides a method for the potentiation of cell mediated immunity against persistent viral infections by administration to a host of at least one antiestrogenic agent capable of enhancing
10 cytotoxic T lymphocytes (CTL), which is expected to lead to the elimination of all infected cells and to the complete clearance of the virus and virus induced tumor cells from the body, leaving behind protective cell mediated immunity. The prerequisite for this adjuvant
15 therapy is the presence of MHC restricted virus specific CTL which kill virus infected target cells very specifically. The elimination of virus producing cells leads to virus clearance and lasting cell-mediated immunity. We showed earlier that TX is immunosuppressive
20 (Nagy & Berczi, 1986; Baral et al., 1979). Therefore, TX and possibly TO, have a dual function during therapy with CTL, namely the sensitization of target cells and the inhibition of switch to humoral, nonprotective immune responses.

25 In accordance with one aspect of the invention, immunity in persistently infected hosts can be potentiated to cause viral clearance according to the following steps:

- a. detecting MHC restricted cytotoxic T
30 lymphocytes in the host,
- b. demonstrating *in vitro* that antiestrogens enhance the killing of virus infected target cells, and
- c. treating the infected host with an effective amount of at least one triphenylethylene antiestrogen to
35 potentiate the killing of virus infected cells in a synergistic manner by CTL.

To fortify the immune response in hosts with advanced persistent viral infections in order to induce viral clearance and cure, the present invention, in another aspect, provides a method, as follows:

5 a. expanding killer cells in tissue culture and reinfusing to the host,

b. testing that antiestrogens enhance the cytotoxic action of these cells on virus infected targets, and

10 c. treating the host with an effective amount of at least one antiestrogen.

The enhancement of viral immunity in chronically infected hosts may be effected, in accordance with a further aspect of the invention as follows:

15 a. treating the host with an effective amount of at least one cytokine, such as interleukin-2 (IL-2) or interleukin-12 (IL-12), to induce and/or enhance cell mediated immune responses in the infected host,

b. demonstrating CTL activity and antiestrogen
20 induced enhancement of virus infected target cell destruction, and

c. applying antiestrogen therapy in order to potentiate the killing effect of cytotoxic cells induced.

The treatment of hosts infected with persistent
25 viruses may additionally be carried out as follows:

a. inducing virus specific CTL in the host by vaccination,

b. demonstrating CTL activity and antiestrogen induced enhancement of cytotoxicity, and

30 c. initiating antiestrogen therapy as soon as virus specific CTL is demonstrable in the immunized host.

Interferon- α (IFN α) is being used currently for the treatment of viral hepatitis and some other viral diseases. In order to take advantage of the anti-viral
35 effect of interferon and the enhancement of host immunity by antiestrogens, we tested whether or not interferons

and antiestrogens can be applied jointly. As shown in Tables 3, 4 and 5 below, antiestrogens had a modest enhancing effect on the cytotoxic action of killer cells as did interferons. Joint application of interferons with anti-estrogens resulted in a further increase in cytotoxicity. Interferons are known to induce prostaglandins, which in turn can inhibit cytotoxicity. For this reason, we tested indomethacin (INDO) in order to see whether or not prostaglandin mediated inhibition exists in our system (Rossol et al., 1992). As shown in Tables 4 and 5 below, the joint application of antiestrogen plus interferon- γ plus indomethacin resulted in the highest cytotoxicity. This was true regardless of whether the killer cells were pretreated or when the drugs were applied during the cytotoxic reactions so that both target and effector cells were exposed to them. On the basis of these results, the following recommendations are made for treatment:

a. Use antiestrogen and interferon jointly to enhance host immunity and virus clearance in all situations listed above.

b. Additional treatment with indomethacin, especially in combination with interferon- γ and antiestrogen, is recommended for best results.

DETAILED DESCRIPTION OF THE INVENTION

A. General treatment protocols

1. Treatment of virus infected hosts in an early stage.

a. Virus infection is determined by the demonstration of viremia with the aid of tests for viral nucleic acid and/or viral antigen. An alternate method may be the detection of virus specific antibodies in the serum.

b. The presence in the infected host of major histocompatibility antigen restricted cytotoxic T

BAD ORIGINAL

lymphocytes (MHCR-CTL) using in vitro methods of cytotoxicity is determined. If killer cell activity is found, the effect of TX and TO and other effective antiestrogens on the in vitro cytotoxic reaction is examined.

5 c. If MHCR-CTL is present in the host in sufficient level and TX and/or TO or other antiestrogen is found to sensitize virus infected target cells in vitro for immune cytolysis, the host is treated with at least one antiestrogen. Treatment with antiestrogen may be combined with treatment with at least one interferon and, optionally, indomethacin and maintained until virus clearance is achieved.

10 d. If no significant decrease in viral titer is observed in the blood during four weeks of treatment, additional treatment with interleukin-2 or interleukin-12, which are known to stimulate CTL activity, may be effected.

e. An alternate solution to the problem of insufficient viral clearance is the expansion of MHCR-CTL in tissue culture and reinfusion of the expanded CTL population to the host. Such treatment may be combined with cytokine treatment, such as with IL-2 and/or IL-12.

20 f. Vaccination may be an additional way to increase the level of MHCR-CTL in the host which could then be followed by treatment with antiestrogens in order to promote the destruction of infected cells and viral clearance.

2. Treatment of hosts with advanced infections but no apparent disease.

30 a. The diagnosis of infection and the presence of MHCR-CTL is done as described under 1. In the present case, it is expected that antibodies are present and nonspecific cytotoxic mechanisms also operate (for example, nonspecific CD8⁺ CTL, antibody dependent cellular cytotoxicity, and complement fixing antibodies).

BAD ORIGINAL

whereas MHCR-CTL may be weakly cytotoxic or completely paralyzed. Therapy is contraindicated with central nervous system involvement.

5 b. Reactivate MHCR-CTL in vitro using antigen presenting cells and cytokines, such as IL-2 and/or IL-12.

 c. Establish that antiestrogens enhance cytotoxicity.

10 d. Expand the cultures and reinfuse into the host while antiestrogen treatment is applied.

 e. Additional treatment with IL-2 and/or IL-12 in order to help maintain cytotoxicity at a significant level and treatment with interferons is recommended to limit virus multiplication.

15 3. Vaccination.

 a. The difficulty of vaccination against persistent viral infections is that nonprotective or even enhancing immune reactions may also be induced, which would interfere with MHCR-CTL that mediate host response.
20 Live virus vaccines that are capable of infecting the susceptible cells have been found to be far superior for vaccination than killed viruses. The apparent explanation for this is that only virus infected cells are able to present processed antigen associated with
25 MHC-I which stimulate CTL specifically. Once a sufficient level of MHCR-CTL has been reached after the injection of the vaccine, and enhancement of cytotoxicity by antiestrogens established, antiestrogen treatment may be initiated in order to prevent the switch to humoral
30 immune reactions. The vaccination of infected animals in combination with antiestrogen treatment is expected to lead to viral clearance and complete recovery which may not happen if the vaccine is given alone.

B. Methods.

35 We use HIV infection and AIDS as an example for our methodology. However, it will be apparent to those

BAD ORIGINAL

skilled in the art that the methodology employed herein equally applies to other viral infections, including those described in the Background to the Invention.

1. **Diagnosis of infection.** Quantitative
5 polymerase chain reaction for the detection of proviral DNA of HIV, a reliable assay for the detection of p24 viral antigen in plasma and assays for the detection of virus specific antibodies for routine tests are available (Wood et al., 1993). The DNA-PCR test is most useful for
10 the follow-up of viral burden in patients.

2. **Detection of cytotoxic T lymphocytes.** MHCR-CTL can be assayed by using EBV-transformed B lymphocytes as targets that have been infected with a vaccinia virus carrying HIV genes. Both autologous and allogeneic B
15 cells can readily be infected by such virus and by this means it is possible to test whether or not the CTL detected is MHC restricted (Lamhamedi-Cherradi et al., 1992; Grant et al., 1992; Ho et al., 1993). This test is also suitable for the determination of MHC restriction
20 which is of utmost importance for our treatment protocol.

3. **Drug treatment.** For in vitro assays it is sufficient to treat target cells with TX (up to 1 μ M) and TO (up to 5 μ M) for 4 hours which is followed by labelling with ^{51}Cr and using the target cells in
25 cytotoxicity assays. The most important virus carrying target in HIV infections is the CD4⁺ helper T lymphocyte, although monocyte-macrophages, astrocytes, neurons and epithelial cells may also be infected. The enhancing effect of antiestrogens on the lysis of autologous HIV
30 infected targets by MHCR-CTL is to be established prior to the initiation of treatment.

For the treatment of patients, the doses of TX and TO that are currently in use for cancer therapy may be applied. For TX, a suitable oral dose is about 20 to 40
35 mg given daily (Anderson et al., 1991). The therapeutic dose for TO is about 40 to 60 mg, but can be raised up to

BAD ORIGINAL

about 240 mg daily (Hietanen et al., 1990). Antiestrogen treatment is to be started as soon as MHCR-CTL is present in the patient (for example, endogenous killer cells, vaccine induced killer cells or in vitro activated killer cells reinfused to the patient). The antiestrogen therapy and killer cell therapy (if given) are then maintained jointly, optimally until the complete clearance of the virus. Additional treatment with interferon would help to control virus production and enhance further host immunity.

4. Treatment with cytokines. Interleukin-2 has long been demonstrated to have a therapeutic effect on syngeneic animal tumors and some beneficial effects could also be demonstrated in certain human malignancies. Recently treatment of AIDS patients with IL-2 also produced beneficial effects, especially when used in combination with zidovudine (Schwartz et al., 1991; Mazza et al., 1992; McMahon et al., 1994). $0.2-12 \times 10^6$ IU/sq m doses were used and well tolerated when injected s.c.

Our survey of the cancer literature indicates that the dose of IL-2 which is well tolerated ranges from about 2 to 8 million IU (about 7.5 to 30,000 IU/kg) which is given subcutaneously (s.c) for about 4 to 5 days (often combined with killer cells). Fever, chills, skin rash, anxiety and dizziness may occur as complications of subcutaneous IL-2 treatment (Schoof et al., 1988; Mitchell et al., 1988; Topalian et al., 1988; Hancock, 1991; Sosman et al., 1988). We recommend treatment with 5 million IU/day s.c., which is maintained for 10 days while the patient is also treated with TX or TO or other antiestrogen. This treatment may be repeated several times in a week to 10 day intervals if it is deemed to be necessary while drug treatment is maintained continuously.

Interleukin-12 has been recognized recently as a cytokine with a powerful stimulatory effect of cell

BAD ORIGINAL

mediated immunity, especially of CTL and NK cells. IL-12 has suppressed tumor growth in animal model systems and exerted antimicrobial activity against bacterial, yeast, parasitic and viral models of infection (Chehimi & Trinchieri, 1994; Wolf et al., 1994). IL-12 has also been shown to stimulate CTL responses to viral antigens such as influenza virus and HIV *in vitro*, and is capable of reactivating anergic CTL from HIV infected patients during *in vitro* stimulation (Clerici et al., 1993).

Currently, there is insufficient human experience with IL-12 treatment to recommend a schedule for dosage. It is known, however, that IL-12 is less toxic than is IL-2 so it is to be expected that the dosage would be more lenient. Once a suitable dose has been established, we would recommend to use an effective daily dosage for 10 day intervals jointly with TX or TO treatment similar to our recommendation for IL-2.

The treatment of chronic hepatitis C in patients with interferon- α is of current interest. The standard dose of IFN α 2b is 3 million units/week for 24 weeks. High doses (15 million units or 22.5-30 million units/week) for 24 weeks have also been tested. Higher doses of IFN ameliorated the severity of hepatitis in patients who did not respond to the standard dose; however, a sustained response was not produced as a rule. Side effects were increased with high doses (Bonkovsky et al., 1996). AIDS patients were treated with low dose IFN α (3 million units s.c., 3 x week). This dose was well tolerated and had a beneficial effect on HIV-related thrombocytopenic purpura (Northfelt et al., 1995). Interferon- γ has been used experimentally on cancer patients at doses of 150-200 μ g s.c. (Lummen et al., 1996; Fiehn et al., 1995).

5. Expansion of CTL in culture. Peripheral blood lymphocytes are stimulated with autologous B lymphoblastoid cell lines derived by infection with the

~~BAD ORIGINAL~~

Epstein-Barr virus followed by infection with recombinant vaccinia virus containing HIV genes. Such stimulator cells can be fixed with paraformaldehyde and used as antigen presenting cells to stimulate peripheral blood mononuclear cells from asymptomatic HIV seropositive individuals. Specific cytotoxic activity can be determined by ^{51}Cr release from viable antigen bearing B cell lines used for stimulation and from similar targets generated with B cell lines of unrelated subjects. Interleukin-2 is useful to aid expansion of CTL in this system. HIV specific CTL may be maintained for as long as 11 months and may be reinfused to the donor patients without any significant complications (Walker et al., 1989; Grant et al., 1992; Lamhamedi-Cherradi et al., 1992; Ahmad et al., 1993; Torpey et al., 1993; van Baalen et al., 1993; Whiteside et al., 1993).

6. Adaptive immunotherapy of AIDS patients with in vitro expanded MHC-CTL. There is not enough experience with regard to the adaptive immunotherapy of AIDS patients with in vitro generated CTL for the establishment of a definite treatment schedule. However, in cancer immunotherapy there is somewhat more experience and a cell dose of about 1×10^8 to 1×10^{10} has been suggested as being effective for treatment with lymphokine activated killer cells. The cells may be infused every day or every other day for up to about 10 days. The patient may require about 1 to 4 such treatments sequentially before achieving tumor remission. CTL may be administered in a similar fashion (Rosenberg et al., 1987; Fujimoto, 1992; Horvath et al., 1993). We recommend daily intravenous infusion with about 5×10^8 to 10^{10} MHC-CTL for 5 days, which is to be repeated in 10 day intervals, while the patient is maintained on TX or TO continuously. Additional treatment with interferon is desirable.

BAD ORIGINAL

7. Vaccination. Hu et al. (1989) vaccinated macaques with recombinant vaccinia virus which expressed the envelope glycoprotein or simian type D retrovirus. The vaccinated animals showed complete protection when challenged with pathogenic doses of simian AIDS virus. HIV infected subjects have been vaccinated similarly with recombinant vaccinia virus expressing HIV glycoprotein 160. Such vaccination resulted in MHC restricted cytotoxic T lymphocytes showing specificity for gp160 (Kundu et al., 1992; el-Daher et al., 1993). In the anticipation that effective vaccine(s) will be developed against HIV in the future, we recommend the treatment schedule outlined below.

Once viral immunity (mediated by MHC-CTL) has been induced in HIV infected patients, we recommend immediately the initiation of TX or TO treatment as described above. Additional treatment with IL-2 or IL-12 is also recommended in order to support the long term maintenance of virus specific MHC restricted CTL. If the level of CTL induced is judged to be unsatisfactory, one may consider the in vitro expansion of CTL and reinfusion as discussed above. Interferon treatment should be considered as an adjunct therapy, especially if the viral load is significant.

EXAMPLES

A series of experiments were carried out, the results of which are outlined in Tables 1 to 5 below.

Example 1:

This Example shows the effect of tamoxifen (TX) and toremifene (TO) on lysis of Epstein-Barr virus (EBV) by autologous cytotoxic T lymphocytes.

In a first set of experiments, cytotoxic lymphocytes were generated by co-culturing Epstein-Barr virus (EBV) transformed B lymphocytes with autologous peripheral blood mononuclear cells for 5 days or longer. Tamoxifen (TX) (1 μ M) or toremifene (TO) (5 μ M) treatment of the

BAD ORIGINAL

target cells was effected for 4 hours, prior to labelling the target cells with ^{51}Cr . The cytotoxicity reaction was carried out at a 1:25 target:effector ratio for four hours.

5 The results of this first set of experiments is set forth in Table 1 below. As may be seen from this Table, TX and TO potentiated the lysis of EBV-transformed B lymphocytes by autologous cytotoxic T lymphocytes.

Example 2:

10 This Example shows the effects of TX and TO on lysis of Epstein-Barr virus (EBV) by autologous LAK cells.

In a second set of experiments, lymphokine activated killer (LAK) cells were generated by the stimulation of autologous peripheral blood lymphocytes ($2 \times 10^6/\text{ml}$) with human recombinant interleukin-2 (500 IU/ml) for 6 days. TX ($1 \mu\text{M}$) and TO ($5 \mu\text{M}$) treatment of target cells was effected for 4 hours prior to labelling the cells with ^{51}Cr . The cytotoxicity reaction (1:25 effector:target ratio) was terminated at 4 hours.

20 The results of this second set of experiments are set forth in Table 2 below. As may be seen from Table 2, TX and TO potentiated the lysis of EBV-transformed B lymphocytes by autologous LAK cells.

Example 3:

25 This Example shows the effect of interferon on the cytotoxic action of TX and TO.

Next we tested whether or not treatment of the cytotoxic cells with TX, TO, or with IFN α would affect their cytotoxic action. The treatment shown in Table 3 was applied to CTL for 4 hours (TX at $1 \mu\text{M}$, TO at $5 \mu\text{M}$, and IFN α at 500 IU/ml), which was followed by cytotoxicity assay (^{51}Cr release) for another 4 hours at 1:25 target:effector ratios. As is shown in Table 3, interferon did increase the cytotoxic action of CTL as did TX and TO, and treatment with both agents resulted in the highest cytotoxicity.

BAD ORIGINAL

Example 4:

This Example shows the effect of interferon- α , interferon- γ and indomethacin in combination with TO on target cells.

5 In the next series of experiments, we extended our studies for the treatment of target cells with interferon- α , interferon- γ (100 IU/ml), and indomethacin (5 μ g/ml), in combination with TO (5 μ M). The results are given in Table 4. Experiments with non-treated
10 targets indicate that the treatment of effector cells with TO, IFN- α plus INDO raised significantly the mean cytotoxicity from $9.6 \pm 1.5\%$ to $22.5 \pm 0.9\%$ ($P < 0.01$). When cytotoxicity on non-treated targets was compared to TO treated targets, significant enhancement occurred with
15 CTL after TO treatment ($P < 0.05$), after IFN- γ plus INDO treatment ($P < 0.05$) and with CTL treated with TO plus IFN- γ plus INDO ($P < 0.01$). These experiments indicate that the enhancement of cytotoxicity by TO will take place in the presence of interferon, and that interferon- γ
20 especially in combination with indomethacin actually enhances significantly the cytotoxic action of killer cells (Table 4).

Example 5:

This Example shows the effect of IFN- α , IFN- γ , TO
25 and INDO and combinations thereof.

Finally, we exposed both the target and the effector cells to IFN- α , IFN- γ , TO and INDO, and to their various combinations (at concentrations given for Table 5) for the duration of the cytotoxic interaction (6 or 18 hrs).
30 In case of 6 hr treatment a significant elevation of cytotoxicity was observed with TO treatment alone, and TO plus INDO ($P < 0.05$), IFN- α plus TO plus INDO and IFN- γ plus TO plus INDO ($P < 0.01$). However, no further enhancement of the cytotoxic reaction was achieved by the
35 addition of IFN and INDO compared to TO alone. The trend obtained in 18 hour cultures was similar. Here target cell destruction reached its highest value in the

BAD ORIGINAL

cultures treated with TO plus IFN- γ plus INDO ($P < 0.01$). These experiments indicate that both the antiestrogens (in this case, TO) can enhance the cytotoxic effect of CTL and that additional treatment with interferons and
5 INDO does not interfere with this effect. Therefore, it seems feasible that one can enhance the destruction of virus infected target cells by antiestrogens while interfering with virus production using interferon- α or - γ . Taken together, these experiments indicate that the
10 treatment of viral diseases, such as HIV infection or hepatitis C, with interferons can be improved if additional treatment with antiestrogens is applied. Indomethacin can be used to oppose the generation of immunosuppressive prostaglandin E by interferon
15 treatment.

SUMMARY OF DISCLOSURE

In summary of this disclosure, a method is provided for the potentiation of cell-mediated immunity in an
20 infected host against persistent viral infection by the administering to the host, which may be a human or animal host, an effective amount of at least one antiestrogenic agent, which may be a triphenylethylene antiestrogen, including tamoxifen and toremifene. This treatment is
25 compatible with interferon therapy, which is known to limit the production of viral particles. Combination treatment with antiestrogens and interferon is expected to lead to better results than interferon treatment alone. Modifications are possible within the scope of
30 this invention.

BAD ORIGINAL

TABLE 1

The anti-estrogens, tamoxifen (TX) and toremifene (TO), enhance the killing of EBV infected target cells by autologous cytotoxic T lymphocytes (CTL).

Exp.	Treatment	Percent lysis	Isotope release (^{51}Cr cpm \pm SE)	
			Spontaneous	Total
1	None	3 ± 0.3	189 ± 15	1925 ± 141
	TX	22 ± 1.2^a	179 ± 20	1889 ± 94
	TO	18 ± 0.6^b	190 ± 11	1912 ± 73
52	None	14 ± 1.1	177 ± 12	1657 ± 111
	TX	34 ± 2^b	184 ± 25	1701 ± 98
	TO	32 ± 0.9^b	196 ± 17	1715 ± 135
3	None	4 ± 0	279 ± 12	2703 ± 48
	TX	15 ± 0.6^b	265 ± 11	2561 ± 71
	TO	14 ± 0.3^b	236 ± 17	2447 ± 94
4	None	3 ± 0.3	289 ± 20	2613 ± 25
	TX	12 ± 1.1^a	274 ± 16	2648 ± 125
	TO	11 ± 0.3^a	280 ± 15	2697 ± 98
5	None	3 ± 0	284 ± 7	2645 ± 145
	TX	14 ± 1.1^b	275 ± 11	2579 ± 94
	TO	12 ± 0.3^b	291 ± 13	2701 ± 111

Statistics (t test): Controls with no treatment were compared with groups treated with TX or TO.

a = $P < 0.05$; b = $P < 0.01$; c = $P < 0.001$

BAD ORIGINAL

TABLE 2

The antiestrogens, TX and TO, enhance the killing of EBV infected target cells by autologous lymphokine activated killer (LAK) cells.

Exp	Treatment	Percent lysis	Isotope release (^51Cr , cpm \pm SE)	
			Spontaneous	Total
1	None	1 ± 0	257 ± 15	1678 ± 114
	TX	10 ± 0.6^b	284 ± 19	1934 ± 198
	TO	16 ± 1.5^c	361 ± 27	2067 ± 157
2	None	3 ± 0.3	238 ± 31	2001 ± 113
	TX	15 ± 1.3^b	274 ± 25	2484 ± 177
	TO	17 ± 0.5^c	297 ± 18	2613 ± 194

Statistics: Please see legends to Table 1.

BAD ORIGINAL

TABLE 1

The effect of TX, TO and IFN α on the cytotoxic action of CTL.

Treatment of CTL	Percent lysis
none	1 \pm 0
IFN α	7 \pm 0.9 ^b
TX	5 \pm 0.7 ^a
TO	8 \pm 0.3 ^b
IFN α + TX	18 \pm 2.9 ^c
IFN α + TO	29 \pm 1.2 ^c

Please see legends to Table 1.

BAD ORIGINAL

TABLE 4

The effect of TO, IFN α , - γ and INDO on the cytotoxic potential of CTL.

Treatment of CTL	Exp:	Percent lysis					
		1	2	Mean \pm SE ¹	3 (TO)	4 (TO)	Mean \pm SE ²
None		6 \pm 0.8	13 \pm 0.6	9.6 \pm 1.5	13 \pm 0.6	23 \pm 1.1	18 \pm 2.3
IFN α		10 \pm 0.6	14 \pm 0.9	12 \pm 1.2	16 \pm 0.7	26 \pm 0.9	21.5 \pm 2.3
IFN γ		11 \pm 0.9	15 \pm 0.3	13 \pm 5.1	18 \pm 1.1	27 \pm 0.3	19.3 \pm 3.1
TO		17 \pm 0.9	14 \pm 0.3	16 \pm 0.8	24 \pm 0.6	29 \pm 1.2	26.5 \pm 1.2
INDO		8 \pm 0.3	13 \pm 1.1	10.6 \pm 1.1	14 \pm 1.1	20 \pm 0.6	17 \pm 1.4
IFN α + INDO		10 \pm 1.1	15 \pm 0.6	12.3 \pm 1.2	17 \pm 0.3	18 \pm 1.1	17.3 \pm 0.6
IFN γ + INDO		13 \pm 0.9	20 \pm 0.6	16.6 \pm 1.5	24 \pm 0.9	30 \pm 1.7	26.8 \pm 1.6
TO + INDO		17 \pm 1.1	16 \pm 0.3	16.6 \pm 0.6	27 \pm 1.5	31 \pm 1.2	25.3 \pm 3.3
IFN α + INDO + TO		19 \pm 1.5	17 \pm 1.1	18 \pm 1	27 \pm 0.3	31 \pm 1.7	24.1 \pm 4
IFN γ + INDO + TO		22 \pm 1.4	23 \pm 1.2	22.5 \pm 0.9 ^b	32 \pm 1.1	39 \pm 0.9	35.6 \pm 1.8 ^b

¹ Mean \pm SE was calculated from exps 1 and 2, which were conducted with non-treated target cells.

² Mean \pm SE was calculated from exps 3 and 4, which were done with TO treated target cells.

Statistics (t test): Mean cytotoxicity of control groups compared to all other groups using the same target cells - untreated vs TO + IFN γ + INDO (P<0.01). Controls compared to all other groups - not significant. Mean cytotoxicity on untreated targets compared to TO treated targets - CTL treated with TO or IFN γ + INDO treatment (P<0.05); CTL treated with TO + IFN γ + INDO (P<0.01).

TABLE 5

The killing potential of CTL is increased by the presence during the cytotoxic reaction of TO, interferon- γ (IFN γ) and indomethacin (INDO).

Treatment	Percent lysis					
	Exp:	1 (6 h)	2 (6 h)	Mean \pm SE ¹	3 (18 h)	4 (18 h)
None		9 \pm 0.7	10 \pm 1.1	9.3 \pm 0.6	17 \pm 1.5	20 \pm 1.1
IFN α		7 \pm 0.3	10 \pm 0.3	8.8 \pm 0.7	16 \pm 0.9	18 \pm 0.3
IFN γ		8 \pm 0.3	11 \pm 1.1	9.6 \pm 0.8	21 \pm 0.6	22 \pm 0.6
TO		20 \pm 0.8	19 \pm 0.3	19.5 \pm 0.6 ¹	25 \pm 2.9	29 \pm 2.6
INDO		8 \pm 1.1	9 \pm 1.1	8.5 \pm 0.7	17 \pm 0.3	17 \pm 1.1
IFN α + INDO		12 \pm 0.6	14 \pm 1.2	13 \pm 0.6	18 \pm 1.4	19 \pm 1.2
IFN γ + INDO		14 \pm 0.3	19 \pm 0.6	16.3 \pm 1.2	18 \pm 1.2	23 \pm 1.4
TO + INDO		18 \pm 0.9	20 \pm 0.6	18.6 \pm 0.9 ¹	22 \pm 1.1	22 \pm 1.1
IFN α + INDO + TO		20 \pm 1.1	20 \pm 1.1	20 \pm 1 ¹	21 \pm 1.5	25 \pm 0.9
IFN γ + INDO + TO		24 \pm 0.3	22 \pm 1.2	23 \pm 0.7 ¹	29 \pm 1.4	33 \pm 1.1

¹ Mean cytotoxicity \pm standard errors was calculated from experiments 1 and 2, which were conducted for 6 hours.

² Mean cytotoxicity \pm standard errors was calculated from experiments 3 and 4, which were conducted for 18 hours.

Statistics (t test): Mean cytotoxicity of control groups with no treatment in the 6 hr experiment compared to all other groups - significant increase in cytotoxicity occurred with TO and TO + INDO ($P < 0.05$), and IFN α + TO + INDO and IFN γ + TO + INDO ($P < 0.01$). Controls compared to all other groups - not significant. In the 18 hr experiment, cytotoxicity is significantly higher than in controls in the IFN γ + TO + INDO treated group ($P < 0.01$). The other groups are not significantly different.

BAD ORIGINAL

REFERENCES

- Ahmad, A., A. Ladha, E.A. Cohen, and J. Menezes. Stable expression of the transfected HIV-1 env gene in a human B cell line: characterization of gp120-expressing clones and immunobiological studies. *Virology* 192:447-457, 1993.
- Andersson, M., H.H. Storm, and H.T. Mouridsen. Incidence of new primary cancers after adjuvant tamoxifen therapy and radiotherapy for early breast cancer. *J. Natl. Cancer Inst.* 83:1013-1017, 1991.
- van Baalen, C.A., M.R. Klein, A.M. Geretti, R.I. Keet, F. Miedema, C.A. van Els, and A.D. Osterhaus. Selective in vitro expansion of HLA class I-restricted HIV-1 Gag-specific CD8+ cells: cytotoxic T-lymphocyte epitopes and precursor frequencies. *AIDS* 7:781-786, 1993.
- Baral, E., S. Kwok, I. Berczi. The influence of estradiol and tamoxifen on the mixed lymphocyte reaction in rats. *Immunopharmacology* 21:191-198, 1991.
- Berczi, I., P. Strausbauch, A.H. Schon. Rejection of tumor cells in vitro. *Science* 180:1289-1291, 1973.
- Bonkovsky, H.L., B.D. Clifford, L.J. Smith, C. Allan, and B. Banner. High-dose interferon-alpha 2b for re-treatment of nonresponders or relapsing patients with chronic hepatitis C. A controlled randomized trial. *Dig. Dis. Sci.* 41:149-154, 1996.
- Chehimi, J., and G. Trinchieri. Interleukin-2: a bridge between innate resistance and adaptive immunity with a role in infection and acquired immunodeficiency. *J. Clin. Immunol.* 14:149-161, 1994.
- Chow, V.T. Cancer and viruses. *Ann. Acad. Med. Singapore* 22:163-169, 1993.
- Clerici, M., D.R. Lucey, J.A. Berzofsky, L.A. Pinto, T.A. Wynn, S.P. Blatt, M.J. Dolan, C.W. Hendrix, S.F. Wolf, and G.M. Shearer. Restoration of HIV-specific cell-mediated immune responses by interleukin-12 in vitro. *Science* 262:1721-1724, 1993.
- el Daher, N., M.C. Keefer, R.C. Reichman, R. Dolin, and N.J. Roberts Jr. Persisting human immunodeficiency virus type 1 gp160-specific human T lymphocyte responses including CD8+ cytotoxic activity after receipt of envelope vaccines. *J. Infect. Dis.* 168:306-313, 1993.

- Daniel, R.C., M.H. Gatei, A.F. Good, D.B. Boyle, and M.F. Lavin. Recombinant viral vaccines for enzootic bovine leucosis. *Immunol. Cell. Biol.* 71:399-404, 1993.
- Fenner, F.J. and D.O. White. *Medical Virology*, 2nd edition, Academic Press, 1976.
- Fiehn, C., O. Prummer, H. Gallati, B. Heilig, and W. Hunstein. Treatment of systemic mastocytosis with interferon-gamma: failure after appearance of anti-IFN-gamma antibodies. *Eur. J. Clin. Invest.* 25:615-618, 1995.
- Fujimoto, S., K. Araki, S. Hamasato, N. Maeda, and Y. Noguchi. Development of specific immunotherapy for cancer. *Hum. Cell.* 5:247, 1992.
- Grant, M.D., F.M. Smaill, D.P. Singal, and K.L. Rosenthal. The influence of lymphocyte counts and disease progression on circulating and inducible anti-HIV-1 cytotoxic T-cell activity in HIV-1-infected subjects. *AIDS* 6:1085-1094, 1992.
- Hancock, G.E., A. Molloy, B.K. Ab, R. Kiessling, M. Becc-Bleumink, Z.A. Cohn, and G. Kaplan. In vivo administration of low-dose human interleukin-2 induces lymphokine activated killer cells for enhanced cytotoxicity in vitro. *Cell. Immunol.* 132:277, 1991.
- Hietanen, T., D. Baltina, R. Johansson, S. Numminen, T. Hakala, L. Helle, and R. Valavaara. High dose toremifene (240 mg daily) is effective as first line hormonal treatment in advanced breast cancer. An ongoing phase II multicenter Finnish-Latvian cooperative study. *Breast Res. Treat.* 16 (Suppl):S37-S40, 1990.
- Ho, H.N., L.E. Hultin, R.T. Mitsuyasu, J.L. Matud, M.A. Hausner, D. Bockstoe, C.C. Chou, S. O'Rourke, J.M. Taylor, and J.V. Giorgi. Circulating HIV-specific CD8+ cytotoxic T cells express CD38 and HLA-DR antigens. *J. Immunol.* 150:3070-3079, 1993.
- Horvath, J., M. Szabo-Szabari, and J.G. Sinkovics. Autologous activated lymphocyte therapy in a community hospital. *Acta Microbiol. Immunol. Hungar.* 41:205, 1994.
- Hu, S.L., J.M. Zarling, J. Chinn, B.M. Travis, P.A. Moran, J. Sias, L. Kuller, W.R. Morton, G. Heidecker, and R.E. Benveniste. Protection of macaques against simian AIDS by immunization with a recombinant vaccinia virus expressing the envelope glycoproteins

- of simian type D retrovirus. *Proc. Natl. Acad. Sci. USA* 86:7213-7217, 1989.
- Ioannides, C.G., B. Fisk, K.R. Jerome, T. Irimura, J.T. Wharton, and O.J. Finn. Cytotoxic T cells from ovarian malignant tumors can recognize polymorphic epithelial mucin core peptides. *J. Immunol.* 151:3693-3703, 1993.
- Kennedy, P.G. Neurological aspects of lentiviral infections in animals. *Baillieres Clin. Neurol.* 1:41-59, 1992.
- Kimelman, D., J.S. Miller, D. Porter, and B.E. Roberts. Ela regions of the human adenoviruses and of the highly oncogenic simian adenovirus 7 are closely related. *J. Virol.* 53:399-409, 1985.
- Koss, L.G. Current concepts of intraepithelial neoplasia in the uterine cervix (CIN). *Appl. Pathol.* 5:7-18, 1987.
- Kradin, R.L., D.S. Lazarus, S.M. Dubinett, J. Gifford, B. Grove, J.T. Kurnick, F.I. Preffer, C.E. Pinto, E. Davidson, R.J. Callahan, and H.W. Strauss. Tumour-infiltrating lymphocytes and interleukin-2 in treatment of advanced cancer. *Lancet* March 18, 577-580, 1989.
- Kundu, S.K., D. Katzenstein, L.E. Moses, and T.C. Merigan. Enhancement of human immunodeficiency virus (HIV)-specific CD4+ and CD8+ cytotoxic T-lymphocyte activities in HIV-infected asymptomatic patients given recombinant gp160 vaccine. *Proc. Natl. Acad. Sci. USA* 89:11204-11208, 1992.
- Lamhamedi-Cherradi, S., B. Culmann-Penciolelli, B. Guy, M.P. Kieny, F. Dreyfus, A.G. Saimot, D. Sereni, D. Sicard, J.P. Levy, and E. Gomard. Qualitative and quantitative analysis of human cytotoxic T-lymphocyte responses to HIV-1 proteins. *AIDS* 6:1249-1258, 1992.
- Lumen, G., M. Goepel, S. Mollhoff, A. Hinke, T. Otto, and H. Rubben. Phase II study of interferon-gamma versus interleukin-2 and interferon-alpha 2b in metastatic renal cell carcinoma. *J. Urol.* 155:455-458, 1996.
- Mazza, P., M. Bocchia, F. Tumietto, P. Costigliola, O. Coronado, G. Bandini, R. Conte, E. Ricchi, N. Vianelli, E. Raine, et al. Recombinant interleukin-2 (rIL-2) in acquired immune deficiency syndrome (AIDS): preliminary report in patients with lymphoma associated with HIV infection. *Eur. J. Haematol.* 49:1-6, 1992.

- McMahon, D.K., J.A. Armstrong, X.L. Huang, C.R. Rinaldo Jr., P. Gupta, T.L. Whiteside, G.J. Pazin, C. Tripoli, and M. Ho. A phase I study of subcutaneous recombinant interleukin-2 in patients with advanced HIV disease while on zidovudine. *AIDS* 8:59-66, 1994.
- Mitchell, M.S., R.A. Kempf, W. Harel, H. Shau, W.D. Boswell, S. Lind, and E.C. Bradley. Effectiveness and tolerability of low-dose cyclophosphamide and low-dose intravenous interleukin-2 in disseminated melanoma. *J. Clin. Oncol.* 4:409, 1988.
- Nagy, E. and I. Berczi. Immunomodulation by tamoxifen and pergolide. *Immunopharmacology* 12:145-153, 1986.
- Northfelt, D.W., E.D. Charlebois, M.I. Mirza, C. Child, L.D. Kaplan, and D.I. Abrams. Continuous low-dose interferon-alpha therapy for HIV-related immune thrombocytopenic purpura. *J. Acquir. Immune Defic. Syndr. Hum. Retrovirol.* 8:45-50, 1995.
- Okazaki, W., H.G. Purchase, and L.B. Crittenden. Pathogenicity of avian leukosis viruses. *Avian Dis.* 26:553-559, 1982.
- Oldstone, M.B., and G.F. Rall. Mechanism and consequence of viral persistence in cells of the immune system and neurons. *Intervirology* 35:116-121, 1993.
- Riddell, S.R., M.J. Gilbert, and P.D. Greenberg. CD8+ cytotoxic T cell therapy of cytomegalovirus and HIV infection. *Curr. Opin. Immunol.* 5:484-491, 1993.
- Rinaldo, C.R. Jr., and D.J. Torpey III. Cell-mediated immunity and immunosuppression in herpes simplex virus infection. *Immunodeficiency* 5:33-90, 1993.
- Roger, P.M., C. Pradier, and P. Dellamonica. Mechanisms of lymphopenia in HIV infection. *Presse Med.* 23:89-94, 1994.
- Rosenberg, S.A., M.T. Lotze, L.M. Muul, A.E. Chang, F.P. Avis, S. Leitman, W.M. Lineham, C.N. Robertson, R.E. Lee, J.T. Rubin, et al. A progress report on the treatment of 157 patients with advanced cancer using lymphokine-activated killer cells and interleukin-2 or high dose interleukin-2 alone. *N. Engl. J. Med.* 316:889, 1987.
- Rossol, S., G. Gianni, R. Rossol-Voth, H. Gallati, W.E. Muller, and K.H. Meyer-zum-Buschenfeld. Cytokine-mediated regulation of monocyte/macrophage cytotoxicity in human immunodeficiency virus-1

BAD ORIGINAL

- infection. *Med. Microbiol. Immunol. Berl.* 181:267-281, 1992.
- Rowland-Jones, S., J. Sutton, K. Ariyoshi, T. Dong, F. Gotch, S. McAdam, D. Whitty, S. Sabally, A. Gallimore, T. Corrah, M. Takiguchi, T. Schultz, A. McMichael, and H. Whittle. HIV-specific cytotoxic T-cells in HIV-exposed but uninfected Gambian women. *Nature Med.* 1:59-64, 1995.
- Schendel, D.J. and B. Gansbacher. Tumor-specific lysis of human renal cell carcinomas by tumor-infiltrating lymphocytes: modulation of recognition through retroviral transduction of tumor cells with interleukin 2 complementary DNA and exogenous α interferon treatment. *Cancer Res.* 53:4020-4025, 1993.
- Schoof, D.D., B.A. Gramolini, D.L. Davidson, A.F. Massaro, R.E. Wilson, and T.J. Eberlein. Adoptive immunotherapy of human cancer using low-dose recombinant interleukin 2 and lymphokine-activated killer cells. *Cancer Res.* 48:5007, 1988.
- Schwartz, D.H., G. Skowron, and T.C. Merigan. Safety and effects of interleukin-2 plus zidovudine in asymptomatic individuals infected with human immunodeficiency virus. *J. Acquir. Immune Defic. Syndr.* 4:11-23, 1991.
- Seligman, M. Immunological features of human immunodeficiency virus disease. *Baillieres Clin. Haematol.* 3:37-63, 1990.
- Sellers, R.F. Blue tongue and related diseases. In: "Virus Diseases of Food Animals." E.P.J. Gibbs (Ed.), Academic Press, 1981, Vol. 2, pp. 566-584.
- Sellon, D.C., F.J. Fuller, and T.C. McGuire. The immunopathogenesis of equine infectious anemia virus. *Virus Res.* 32:111-138, 1994.
- Sosman, J.A., P.C. Kohler, J. Hank, K.H. Moore, R. Bechhofer, B. Storer, and P.M. Sondel. Repetitive weekly cycles of recombinant human interleukin-2: responses of renal carcinoma with acceptable toxicity. *J. Natl. Cancer Inst.* 80:60, 1988.
- Thiel, H.J., G. Meyers, R. Stark, N. Tautz, T. Rumenapf, G. Unger, and K.K. Conzelmann. Molecular characterization of positive strand RNA viruses: pestiviruses and the porcine reproductive and respiratory syndrome virus (PRRSV). *Arch. Virol.* Suppl. 7:41-52, 1993.

- Tishon, A., P. Borrow, C. Evans, and M.B. Oldstone. Virus-induced immunosuppression. 1. Age at infection relates to a selective or generalized defect. *Virology* 195:397-405, 1993.
- Topalian, S.L., D. Solomon, F.P. Avis, A.E. Chang, D.L. Freerksen, W.M. Linehan, M.T. Lotze, C.N. Robertson, C.A. Seipp, P. Simon, C.G. Simpson, and S.A. Rosenberg. Immunotherapy of patients with advanced cancer using tumor-infiltrating lymphocytes and recombinant interleukin-2: a pilot study. *J. Clin. Oncol.* 6:839-853, 1988.
- Torpey, D. III, X.L. Huang, J. Armstrong, M. Ho, T. Whiteside, D. McMahon, G. Pazin, R. Heberman, P. Gupta, C. Tripoli, et al. Effects of adoptive immunotherapy with autologous CD8+ T lymphocytes on immunologic parameters: lymphocyte subsets and cytotoxic activity. *Clin. Immunol. Immunopathol.* 66:263-272, 1993.
- de la Torre, J.C., P. Borrow, and M.B. Oldstone. Viral persistence and disease: cytopathology in the absence of cytolysis. *Br. Med. Bull.* 47:838-851, 1991.
- Walker, B.D., C. Flexner, K. Birch-Limberger, L. Fisher, T.J. Paradis, A. Aldovini, R. Young, B. Moss, and R.T. Schooley. Long-term culture and fine specificity of human cytotoxic T-lymphocyte clones reactive with human immunodeficiency virus type 1. *Proc. Natl. Acad. Sci. USA* 86:9514-9518, 1989.
- Whiteside, T.L., E.M. Elder, D. Moody, J. Armstrong, M. Ho, C. Rinaldo, X. Huang, D. Torpey, P. Gupta, D. McMahon, et al. Generation and characterization of ex vivo propagated autologous CD8+ cells used for adoptive immunotherapy of patients infected with human immunodeficiency virus. *Blood* 81:2085-2092, 1993.
- Wolf, S.F., D. Sieburth, and J. Sypek. Interleukin 12: a key modulator of immune function. *Stem Cells* Dayt. 12:154-168, 1994.
- Wood, R., J.G. Montoya, S.K. Kundu, D.H. Schwartz, and T.C. Merigan. Safety and efficacy of polyethylene glycol-modified interleukin-2 and zidovudine in human immunodeficiency virus type 1 infection: a phase I/II study. *J. Infect. Dis.* 167:519-525, 1993.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A method for the potentiation of cell mediated immunity against persistent viral infection in a host, which comprises:

administering to a host an effective amount of at least one antiestrogen capable of enhancing cytotoxic T lymphocytes (CTLs) in the host.

2. The method of claim 1 wherein said host harbours MHC-restricted CTLs and said at least one antiestrogen is initially tested in vitro to demonstrate enhancement of lysis of virus infected cells.

3. The method of claim 2 wherein the antiestrogen is a triphenylethylene antiestrogen.

4. The method of claim 2 wherein said antiestrogen is tamoxifen or toremifene.

5. The method of claim 2 wherein said at least one antiestrogen is administered along with at least one interferon.

6. The method of claim 5 wherein said antiestrogen is tamoxifen or toremifene and said at least one interferon is interferon- α or interferon- γ .

7. The method of claim 6 wherein said at least one antiestrogen and at said at least one interferon are administered along with indomethacin.

8. The method of claim 1 wherein said host does not harbour MHC-restricted cytotoxic T lymphocytes, and MHC-restricted cytotoxic lymphocytes are activated and expanded in tissue culture and reinfused into the host and treatment with at least one antiestrogen is initiated with the first infusion of CTL and maintained for the entire treatment period.

9. The method of claim 8 wherein the antiestrogen is a triphenylethylene antiestrogen.

10. The method of claim 8 wherein said antiestrogen is tamoxifen or toremifene.
11. The method of claim 8 wherein said at least one antiestrogen is administered along with at least one interferon.
12. The method of claim 11 wherein said antiestrogen is tamoxifen or toremifene and said at least one interferon is interferon- α or interferon- γ .
13. The method of claim 12 wherein said at least one antiestrogen and at said at least one interferon are administered along with indomethacin.
14. The method of claim 1 wherein said host is initially treated with at least one immunostimulatory cytokine to stimulate MHC-restricted CTLs.
15. The method of claim 14 wherein said immunostimulatory cytokine is interleukin-2 or interleukin-12.
16. The method of claim 15 wherein the antiestrogen is a triphenylethylene antiestrogen.
17. The method of claim 15 wherein said antiestrogen is tamoxifen or toremifene.
18. The method of claim 15 wherein said at least one antiestrogen is administered along with at least one interferon.
19. The method of claim 18 wherein said antiestrogen is tamoxifen or toremifene and said at least one interferon is interferon- α or interferon- γ .
20. The method of claim 19 wherein said at least one antiestrogen and at said at least one interferon are administered along with indomethacin.
21. The method of claim 1 wherein said host is virus infected but otherwise immunocompetent, said host is vaccinated to stimulate MHC-restricted CTLs, and said host is tested for the presence of MHC-restricted CTLs prior to administration of said at least one antiestrogen.

BAD ORIGINAL

DARBY & DARBY, P.C.
Attn. FEHLNER, Paul F.
805 Third Avenue
New York, N.Y. 10022-7513
UNITED STATES OF AMERICA

Date: 14/03/2001

